

Final
10-2-94
29 CIT
22749
14 P

- a. INSTITUTION:
University of Utah
309 Park Building
1400 East Second South
Salt Lake City, UT 84112
- b. Prevent and Cure Disuse Bone Loss
- c. Final Technical Report, March 1990 - February 1994
- d. NAG-2435
- e. Principal Investigator:
Webster S.S. Jee, Ph.D.
Division of Radiobiology, Bldg. 586]
University of Utah
Salt Lake City, UT 84112
Tel: (801) 581-6366
FAX: (801) 581-7008
- f. NASA Technical Officer: Sara Arnaud, M.D., NASA
Ames Research Center
Moffat Field, CA

Webster S.S. Jee Ph.D.
Webster S.S. Jee, Ph.D.
Principal Investigator

13 September 1994
Date

(NASA-CR-196845) PREVENT AND CURE
DISUSE BONE LOSS Final Technical
Report, Mar. 1990 - Feb. 1994
(Utah Univ.) 14 p

N95-11594

Unclass

G3/52 0022749

TABLE OF CONTENTS

| | Pages |
|--|-------|
| I. PUBLICATIONS | |
| A. Peer review articles. | 1-3 |
| B. Proceeding articles and reviews | 4 |
| II. OBJECTIVES AND HYPOTHESES | 5 |
| III. SUMMARY OF FINDINGS | 5-6 |
| IV. CONTRIBUTIONS TO A BETTER UNDERSTANDING OF BONE BIOLOGY | 6-9 |
| V. CONCLUDING REMARKS | 9 |
| VI. REPRINTS (enclosed). | |

W.S.S. Jee, Ph. D.
NAG-2435

I. PUBLICATIONS

A. Peer Review.

1. Polig, E. and Jee, W.S.S. 1990. A model of osteon closure in cortical bone. *Calcif. Tissue Int.* 47:261-269..
2. Li, X.J. and Jee, W.S.S., Chow, S.Y. and Woodbury, D.M. 1990. Adaptation of cancellous bone to aging and immobilization in the rat: A single photon absorptiometry and histomorphometry study. *Anat. Rec.* 227:12-24.
3. Mori, S., Jee, W.S.S., Li, X.J., Chan, S. and Kimmel, D.B. 1990. Effects of prostaglandin E₂ on production of new cancellous bone in the axial skeleton of ovariectomized rats. *Bone* 11:103-113.
4. Jee, W.S.S., Li, X.J. 1990. Adaptation of cancellous bone to overloading in the adult rat: a single photon absorptiometry and histomorphometry study. *Anat. Rec.* 227:418-426.
5. Li, X.J., Jee, W.S.S., Li, Y.L. and Patterson-Buckendahl, P. 1990. Transient effects of subcutaneously administered Prostaglandin E₂ on cancellous and cortical bone in young adult dogs. *Bone* 11:353-364.
6. Jee, W.S.S., Mori, S., Li, X.J. and Chan, S. 1990. Prostaglandin E₂ enhances cortical bone mass and activates intracortical bone remodeling in intact and ovariectomized female rats. *Bone* 11:253-266.
7. Li, X.J. and Jee, W.S.S. 1991. Adaptation of diaphyseal structure to aging and decreased mechanical loading in the adult rat: A densitometric and histomorphometric study. *Anat. Rec.* 229:291-297.
8. Jee, W.S.S., Li, X.J., Schaffler, M.B. 1991. Adaptation of diaphyseal structure with aging and increased mechanical usage in the adult rat: A histomorphometrical and biomechanical study. *Anat. Rec.* 230:332-338.
9. Jee, W.S.S., Li, X.J. and Ke, H.Z. 1991. The skeletal adaptation to mechanical usage in the rat. *Cells and Materials, Suppl.* 1:131-142.
10. Li, X.J., Jee, W.S.S., Ke, H.Z., Mori, S., Akamine, T. 1991. Age-related changes of cancellous and cortical bone histomorphometry in female Sprague-Dawley rats. *Cells and Materials, Suppl.* 1:25-35.
11. Ke, H.Z., Jee, W.S.S. and Li, X.J. 1991. Partial loss of anabolic effect of prostaglandin E₂ on bone after its withdrawal in rats. *Bone* 12:173-183.
12. Jee, W.S.S., Ke, H.Z. and Li, X.J. 1991. Long term anabolic effects of prostaglandin E₂ on tibial diaphyseal bone in male rats. *Bone and Mineral* 15:33-35.

W.S.S. Jee, Ph.D.
NAG-2435

13. Mori, S., Jee, W.S.S. and Li, X.J. 1992. Production of new trabecular bone in osteopenic ovariectomized rats. *Calcif. Tissue Int.* 50:80-87.
14. Ke, H.Z., Jee, W.S.S., Mori, S., Li, X.J. and Kimmel, D.B. 1992. Effects of long term daily administration of prostaglandin E₂ on maintaining elevated proximal tibial metaphyseal cancellous bone mass in male rats. *Calcif. Tissue Int.* 50:245-252.
15. Akamine, T., Jee, W.S.S., Ke, H.Z., Li, X.J., Lin, B.Y. 1992. PGE₂ prevents disuse-induced cancellous bone loss and adds extra bone to immobilized bones. *Bone* 13:11-22.
16. Jee, W.S.S., Ke, H.Z., Li, X.J. 1992. Loss of PGE₂ induced cortical bone after its withdrawal in rats. *Bone and Mineral*, 17: 31-47.
17. Jee, W.S.S., Akamine, T., Ke, H.Z., Li, X.J., Tang, L.Y. Qeng, Q.Q. 1992. PGE₂ prevents disuse-induced cortical bone loss. *Bone* 13:153-159.
18. Frost, H.M. and Jee, W.S.S. 1992. On the rat model of human osteopenias and osteoporoses. *Bone and Mineral*, 18: 227-236.
19. Jee, W.S.S., Frost, H.M. 1992. Skeletal Adaptation During Growth. *Triangle*. 31:77-88.
20. Ke, H.Z., Jee, W.S.S. 1992. Effects of daily administration of PGE₂ and its withdrawal on the lumbar vertebral bodies in male rats. *Anat. Rec.* 234:172-182.
21. Chen, M.M., Jee, W.S.S., Ke, H.Z., Lin, B.Y., Li, Z.N. and Li, X.J. 1992. Adaptation of cancellous bone to aging and immobilization in the growing rats. *Anat. Rec.* 234: 317-334.
22. Katz, I.A., Jee, W.S.S., Joffe, I.I., Stein, B., Takizawa, M., Jacobs, T.W., Setterberg, R., Lin, B.Y., Tang, L.Y., Ke, H.Z., Zeng, Q.Q., Berlin, J., Epstein, S. 1992. Prostaglandin E₂ alleviates cyclosporin A-induced bone loss in the rat. *J. Bone and Min. Res.* 7: 1191-1200.
23. Tang, L.Y., Jee, W.S.S., Ke, H.Z., Kimmel, D.B. 1992. Restoring and maintaining bone in osteopenic female rat skeleton. I. Changes in Bone Mass and Structure. *J. Bone and Min. Res.* 7:1093-1104.
24. Ke, H.Z., Li, M., Jee, W.S.S. 1992. Prostaglandin E₂ prevents ovariectomy-induced cancellous bone loss in the rats. *Bone and Mineral*. 19: 45-62.
25. Zeng, Q.Q., Jee W.S.S., Ke, H.Z. and Wechter, W.J. (1993). S-Ketoprofen inhibits tenotomy-induced bone loss and dynamics in weaning rats, *Bone and Mineral*. 21:203-218.
26. Ke, H.Z., Jee W.S.S., Zeng, Q.Q., Li, M., and Lin, B.Y. (1993). Prostaglandin E₂ increased rat cortical bone mass when administered immediately following ovariectomy. *Bone and Mineral*, 21:189-201.

W.S.S. Jee, Ph.D.
NAG-2435

27. Ito, H., Ke, H.Z., Jee, W.S.S., and Sakou, T. (1993). Anabolic responses of an adult cancellous bone site to prostaglandin E₂ in the rat,. Bone and Mineral, 21:219-236.
28. Ke, H.Z., Jee, W.S.S., Ito, H., Setterberg, R.B., Li, M., Lin, B.Y., Liang, X.G. and Ma, Y.F. (1993). Greater bone formation induction occurred in aged than young cancellous bone sites. Bone, 14:481-486.
29. Li, M., Jee, W.S.S., Ke, H.Z., Liang, X.G., Lin, B.Y., Ma, Y.F., and Setterberg, R.B. (1993). Prostaglandin E₂ restores cancellous bone to immobilized limb and adds bone to overloaded limb in right hindlimb immobilization rats. Bone, 14:283-288.
30. Jee, W.S.S., Tang, L., Ke, H.Z., Setterberg, R.B., and Kimmel, D.B. (1993). Maintaining restored bone with bisphosphonate in the ovariectomized rat skeleton: Dynamic histomorphometry of changes in bone mass. Bone, 14:493-498.
31. Ma, Y.F., Ke, H.Z., Jee, W.S.S., (1994) Prostaglandin E₂ adds bone to a cancellous bone site with a closed growth plate and low bone turnovers in ovariectomized rats. Bone 15:137-146.
32. Lin, B.Y., Jee, W.S.S., Chen, M.M., Ma, Y.F., Ke, H.Z., Li, X.J. 1994 Mechanical loading modifies ovariectomy-induced cancellous bone loss. Bone & Mineral. 25:199-210.
33. Lin, B.Y., Jee, W.S.S., Ma, Y.F., Ke, H.Z., Kimmel, D.B., Li, X.J. (in press) Effects of prostaglandin E₂ and Risedronate administration on cancellous bone in older rats. Bone.
34. Ma, Y.F., Jee, W.S.S., Ke, H.Z., Lin, B.Y., Liang, X.G., Le, M., Yamamoto, N. (in press) Human parathyroid hormones (1-38) restores cancellous bone to immobilized, osteopenic proximal tibial metaphysis in rats. J.Bone and Mineral.Res.
35. Frost, H.M., Jee, W.S.S. (in press) Perspectives: A vital biomechanical model of endochondral ossification mechanism. Anat. Rec.
36. Frost, H.M., Jee, W.S.S. (in press) Perspectives: Application of biomechanical model of the endochondral ossification mechanism. Anat. Rec.

W.S.S. Jee, Ph.D.
NAG-2435

II. PUBLICATIONS

B. Proceeding articles and Reviews

1. Jee, W.S.S. and X.J. Li. 1990. Small animal models for studying prevention and treatment of osteoporosis. In: Bone Morphometry. H.E. Takahashi (ed.) Nishimura Co, Ltd. Niigata, Japan, pp. 549-554, 1990.
2. Jee, W.S.S. 1990. Augmentation of Bone Mass in Osteoporosis, *ibid* pp. 543-548.
3. Jee, W.S.S. 1990. Local and Systemic Factors Influencing Bone Formation, *ibid* pp. 284-289.
4. Miller, S.C. and Jee, W.S.S. 1990. The bone forming cell: origin, distribution, proliferation and differentiation, *ibid* pp. 29-36.
5. Norrdin, R.W., Jee, W.S.S. and High, W.B. 1990. The role of prostaglandins in bone in vivo. A review. Prostaglandins, leukotrienes and essential fatty acids. 41:139-149.
- 6.. Jee, W.S.S. 1991. Introduction to skeletal function. In: A Basic Science Primer in Orthopaedics. F. Bronner and R.V. Worell (eds.) Williams and Wilkins, pp 3-34.
7. Woodard, J.C. and Jee, W.S.S. 1991 Skeletal System. In: Handbook of Toxicologic Pathology. W.M. Haschek-Holik and C. Rorisseau (eds.), Academic Press, Inc., pp. 489-537.
8. Miller, S.C. and Jee, W.S.S. 1992. Bone lining cells. In: Bone, VOL. 4: Bone Metabolism and Mineralization. B.K. Hall (ed.), Telford Press, Caldwell, NJ, 1-19.
- 9 Jee, W.S.S., Ma, Y.F., Li, M., Liang, X.Q., Lin, B.Y., Li, X.J., Ke, H.Z., Mori, S., Setterberg, R.B., Kimmel, D.B. 1994. Sex steroids and prostaglandin in bone metabolism. In: Sex steroids and bone. Edited by R Ziegler, J. Pfeilschifter and M Brautigam. Ernst Schering Research Foundation Workshop 9. Springer-Verlag, Berlin, pp. 119-150.
10. Jee, W.S.S., Inuoe, J., Jee, K.W., Haba, T., Ke, H.Z., Li, X.J., and Setterberg, R.B. (in press) Histomorphometry assay of growth bones. In: Handbook of Bone Morphometry. Second Edition. H. Takahashi (ed.) Nishimura Co. Ltd. Niigata City, Japan.

W.S.S. Jee, Ph.D.
NAG-2435

II. OBJECTIVES AND HYPOTHESES

- 1) Disuse-related osteopenia and poor trabecular architecture is prevented by non-steroidal anti-inflammatory drugs (NSAID).
- 2) Established disuse-related osteopenia and poor trabecular architecture is cured by PGE₂ or PTH treatment though immobilization continues.
- 3) Though bone mass returns, poor trabecular architecture remains after normal ambulation recovery from immobilization.
- 4) Established disuse-related bone loss and poor trabecular architecture is cured by post-immobilization PGE₂ treatment.

III. SUMMARY OF FINDINGS

1. PGE₂ Treatment Prevented Immobilization-induced Bone Loss.
 - A. PGE₂ administration not only prevented immobilization-induced bone loss, but also added extra bone in a dose-response manner. It stimulated more bone formation than resorption shortened the period of bone remodeling, activated woven bone formation, stimulated lamellar bone formation and shortened the bone resorption and remodeling period to reduce the remodeling space (T. Akamine et al., Prostaglandin E₂ prevents bone loss and adds extra bone to immobilized distal femoral metaphysis in female rats. Bone 13:11-22, 1992)
 - B. PGE₂ treatment-induced increases in periosteal and endocortical bone formation to compensate for the disuse and PGE₂-induced cortical bone loss, and thus prevented immobilization-induced cortical bone loss (W.S.S. Jee et al. Prostaglandin E₂ prevents disuse-induced cortical bone loss. Bone 13:153-159, 1992).
2. S-ketoprofen, a non-steroidal anti-inflammatory drug prevented immobilization-induced bone loss, but not changes in architecture.
 - A. S-ketoprofen, a non-steroidal anti-inflammatory drug, prevented both the inhibition of age-related cancellous bone gain and the stimulation of bone loss induced by immobilization (tenotomy) at the 2.5 mg/kg/d, while it only prevented bone loss at the 0.5 mg/kg/d dose levels. These doses partially protected from the development of poor architecture in that trabecular numbers were reduced.
 - B. In the tibial shaft, S-ketoprofen prevented the immobilization-induced increase in endocortical bone resorption (Q.Q. Zeng et al. S.-Ketoprofen inhibits tenotomy-induced bone loss and dynamics in weanling rats. Bone and Mineral 21:204-218, 1993).

W.S.S. Jee, Ph.D.
NAG-2435

3. PGE₂ restored bone and structure to immobilization-induced osteopenic rats.
Daily 3 or 6 mg/kg/d treatment restores and maintains proximal tibial metaphyseal cancellous bone mass and structure in continuously immobilized tibiae in female rats in spite of continuous immobilization (M. Li et al. Prostaglandin E₂ restores cancellous bone to immobilized limb and adds bone to overloaded limb in right hindlimb immobilization rats. Bone 14:283-288, 1993).
 4. Parathyroid hormone (PTH) restores and adds extra cancellous bone and improves bone structure to immobilized, osteopenic proximal tibial metaphysis in rats.
PTH treatment stimulates cancellous bone formation, restores and adds extra cancellous bone and improves bone structure above that of age-related controls in established, disuse-osteopenic proximal tibial metaphysis of continuously right hindlimb immobilized female rats (Y.F. Ma et al. Human parathyroid hormone (1-38) restores cancellous bone to the immobilized, osteopenic tibial metaphysis in rats. J. Bone and Mineral Research (in press)).
 5. Remobilization after short term immobilization (2 weeks) allows normal bone mass and architecture to return while longer immobilization (> 10 weeks) does not in rats.
(X.G. Liang et al. Effects of remobilization on the immobilization-induced osteopenic rat skeleton. A time response study. In preparation)
- IV. CONTRIBUTIONS TO A BETTER UNDERSTANDING OF BONE BIOLOGY.
1. We found that a powerful anabolic agent like PGE₂ will stimulate direct bone formation during bone modeling and remodeling (1-18).
 2. We found that PGE₂ shortens the phases of the bone remodeling cycle. It reduces the residence time of the remodeling space and enables the agent to activate new bone turnover sites with a smaller transient bone loss phase that could place an individual at increased fracture risk (1-18)
 3. We confirm that bone tissue precedes bone marrow in ontogeny (11,19,20).
 4. Our studies confirm that cortical bone contains a rich source of osteoblastic progenitors, believed to be the lining cells of the cortical bone vascular channels (2,3,4,8,11,13,15).
 5. We found bone mass increases caused by anabolic agents do not continue indefinitely. One might suggest that they plateau at a new level commensurating with the properties of the agent, dose, and the role of skeletal adaptation to mechanical usage (4,6).

W.S.S. Jee, Ph.D.
NAG-2435

6. Our studies indicate that excess bone induced by an anabolic agent disappears after treatment stops. We postulate that bone adapts to mechanical usage by ridding itself of "extra" bone that keeps strain "too low" or adding "extra" bone when strain is "too high" to maintain constant strain within the physiologic window (9,11,17,19,20).
7. In our studies we discovered that we could outsmart the skeleton by using the lose, restore and maintain (LRM) concept to restore and maintain PGE₂ added cancellous and supplement and maintain (SM) concept to build up bone (11,17).
8. We found PGE₂ will stimulate periosteal bone formation and consequently increase the diameter of the cortex and its strength. The engineers tell us that the strength of a given mass can be increased by increasing the diameters of the shaft while making its wall proportionally thinner (2-4,8,13).
9. We found that PGE₂ depresses longitudinal bone growth in male but stimulates it in female immobilized rats. This is an area of research that warrants more study (21).
10. We confirm that the distal tibial metaphysis (DTM) closes at 3 months and a few months later contains a metaphysis with fewer but thicker trabeculae and lower turnover. Studies of this adult metaphysis could improve our understanding of the responses of this adult cancellous bone as well as a low turnover site to bone-seeking agents inexpensively (14,15,18).
11. We discovered that ovariectomy does not but immobilization does induce bone loss in the adult distal tibial metaphyses (18,22).

REFERENCES

1. Mori, S, Jee, WSS., Li, XJ, Chan, S, Kimmel, DB. 1990 Effects of prostaglandin E₂ on production of new cancellous bone in the axial skeleton of ovariectomized rats. *Bone* 11:103-113.
2. Li, XJ, Jee, WSS, Li, YL, Patterson-Buckendahl, P. 1990 Transient effects of subcutaneously administered prostaglandin E₂ on cancellous and cortical bone in young adult dogs. *Bone* 11:353-364.
3. Jee, WSS, Mori, S, Li, XJ, Chan, S. 1990 Prostaglandin E₂ enhances cortical bone mass and activates intracortical bone remodeling in intact and ovariectomized female rats. *Bone* 11:253-266

W.S.S. Jee, Ph.D.
NAG-2435

4. Jee, WSS, Ke, HZ, Li, XJ. 1991 Long term anabolic effects of prostaglandin E₂ on tibial diaphyseal bone in male rats. *Bone and Mineral* 15:33-35.
5. Mori, S, Jee, WSS, Li, XJ. 1992 Production of new trabecular bone in osteopenic ovariectomized rats. *Calcif. Tissue Int.* 50:80-87.
6. Ke, HZ, Jee, WSS, Mori, S, Li, XJ, Kimmel, DB. 1992 Effects of long term daily administration of prostaglandin E₂ on maintaining elevated proximal tibial metaphyseal cancellous bone mass in male rats. *Calcif. Tissue Int.* 50:245-252.
7. Akamine, T, Jee, WSS, Ke, HZ, Li, XJ, Lin, BY. 1992 PGE₂ prevents disuse-induced cancellous bone loss and adds extra bone to immobilized bones. *Bone* 13:11-22.
8. Jee, WSS, Akamine, T, Ke, HZ, Li, XJ, Tang, LY, Qeng, QQ. 1992 PGE₂ prevents disuse-induced cortical bone loss. *Bone* 13:153-159.
9. Ke, HZ, Jee, WSS. 1992 Effects of daily administration of PGE₂ and its withdrawal on the lumbar vertebral bodies in male rats. *Anat. Rec.* 234:172-182.
10. Katz, IA, Jee, WSS, Joffe, II, Stein, B, Takizawa, M, Jacobs, TW, Setterberg, R, Lin, BY, Tang, LY, Ke, HZ, Zeng, QQ, Berlin, J, Epstein S. 1992 Prostaglandin E₂ alleviates cyclosporin A-induced bone loss in the rat. *J. Bone and Min. Res.* 7:1191-1200.
11. Tang, LY, Jee, WSS, Ke, HZ, Kimmel, DB. 1992 Restoring and maintaining bone in osteopenic female rat skeleton. I. Changes in Bone Mass and Structure. *J. Bone and Min. Res.* 7:1093-1104.
12. Ke, HZ, Li, M, Jee, WSS. 1992 Prostaglandin E₂ prevents ovariectomy-induced cancellous bone loss in the rats. *Bone and Mineral* 19:45-62.
13. Ke, HZ, Jee, WSS, Zeng, QQ, Li, M, Lin, BY. 1993 Prostaglandin E₂ increased rat cortical bone mass when administered immediately following ovariectomy. *Bone and Mineral* 21:189-201.
14. Ito, H, Ke, HZ, Jee, WSS, Sakou T. 1993 Anabolic responses of an adult cancellous bone site to prostaglandin E₂ in the rat. *Bone and Mineral* 21:219-236.
15. Ke, HZ, Jee, WSS, Ito, H, Setterberg, RB, Li, M, Lin, BY, Liang, XG, Ma, YF. 1993 Greater bone formation induction occurred in aged than young cancellous bone sites. *Bone* 14:481-486.
16. Li, M, Jee, WSS, Ke, HZ, Liang, XG, Lin, BY, Ma YF, Setterberg, RB. 1993 Prostaglandin E₂ restores cancellous bone to immobilized limb and adds bone to overloaded limb in right hindlimb immobilization rats. *Bone* 14:283-288.

W.S.S. Jee, Ph.D.
NAG-2435

17. Jee, WSS, Tang, L, Ke, HZ, Setterberg, RB, Kimmel, DB. 1993 Maintaining restored bone with bisphosphonate in the ovariectomized rat skeleton: Dynamic histomorphometry of changes in bone mass. *Bone* 14:493-498.
18. Ma, YF, Ke, HZ, Jee, WSS. 1994 Prostaglandin E₂ adds bone to a cancellous bone site with a closed growth plate and low bone turnovers in ovariectomized rats. *Bone* 15:137-146.
19. Ke, HZ, Jee, WSS, Li, XJ. 1991 Partial loss of anabolic effect of prostaglandin E₂ on bone after its withdrawal in rats. *Bone* 12:173-183.
20. Jee, WSS, Ke, HZ, Li, XJ. 1992 Loss of PGE₂ induced cortical bone after its withdrawal in rats. *Bone and Mineral* 17:31-47.
21. Jee, WSS, Ma, YF, Li, M, Liang, XQ, Lin, BY, Li, XJ, Ke, HZ, Mori, S, Setterberg, RB, Kimmel, DB. 1994 Sex steroids and prostaglandin in bone metabolism. In: *Sex Steroids and Bone*. Edited by R. Ziegler, J. Pfeilschifter and M. Brautigam. Ernst Schering Research Foundation Workshop 9. Springer-Verlag, Berlin, pp. 119-150.
22. Ijiri, K, Ma, YF, Jee, WSS, Liang XG. Skeletal adaptation of a non-growing cancellous bone site to aging and immobilization in the rat (in preparation).

V. CONCLUDING REMARKS.

If remobilization after long-term immobilization is incapable of restoring normal bone mass and structure in humans as seen in rats and dogs, anabolic agents like parathyroid hormone and prostaglandin E-like substances will be needed. It is fortunate that several pharmaceutical companies are developing analogues of these two agents. If successful, these agents will be effective for all types of osteoporoses in man if they are tested in an immobilization-induced bone loss model because immobilization-induced bone loss is more difficult to prevent and cure than estrogen-deficiency-induced osteopenia.

REPRINTS

W.S.S. Jee, Ph.D.
NAG-2435

1. Jee, WSS, Ke, HZ, Li, XJ. 1991 Long-term anabolic effects of prostaglandin-E₂ on tibial diaphyseal bone in male rats. *Bone and Mineral* 15:33-55.
2. Jee, WSS, Li, XJ, Schaffler, MB. 1991 Adaptation of diaphyseal structure with aging and increased mechanical usage in the adult rat: A histomorphometrical and biomechanical study. *Anat. Rec.* 230:332-338.
3. Jee, WSS, Li, XJ, Ke, HZ. 1991 The skeletal adaptation to mechanical usage in the rat. *Cells and Materials Suppl.* 1, pp. 131-142.
4. Ke, HZ, Li, XJ, Jee, W.S.S. 1991 Partial loss of anabolic effect of prostaglandin E₁ on bone after its withdrawal in rats. *Bone* 12:173-183.
5. Li, XJ, Jee, WSS, Ke HZ, Mori, S, Akamine T. 1991 Age-related changes of cancellous and cortical bone histomorphometry in female Sprague-Dawley rats. *Cells and Materials Suppl* 1. pp. 25-35.
6. Akamine, T, Jee, WSS, Ke, HZ, Li, XJ, Lin, BY. 1992 Prostaglandin E₂ prevents bone loss and adds extra bone to immobilized distal femoral metaphysis in female rats. *Bone* 13:11-22.
7. Chen, MM, Jee, WSS, Ke, HZ, Lin, BY, Li, QN, Li, XJ. 1992 Adaptation of cancellous bone to aging and immobilization in growing rats. *Anat . Rec.* 234:317-334.
8. Frost, HM, Jee, WSS. 1992 On the rat model of human osteopenias and osteoporoses. *Bone and Min.* 18:227-236.
9. Jee, WSS, Frost, HM. 1992 Skeletal adaptations during growth. *Triangle* 31:77-88.
10. Jee, WSS, Akamine T, Ke, HZ, Li, XJ, Tang, LY, Zeng QQ. 1992 Prostaglandin E₂ prevents disuse-induced cortical bone loss. *Bone* 13:153-159.
11. Jee, WSS, Ke, HZ, Li, XJ. 1992 Loss of prostaglandin E₂-induced extra cortical bone after its withdrawal in rats. *Bone and Mineral* 17:31-47.
12. Katz, IA, Jee, WSS, Joffe, II, Stein, B, Takizawa, M, Jacobs, TW, Setterberg, R, Lin, BY, Tang, LY, Ke, HZ, Zeng, QQ, Berlin, J, Epstein, S. 1992 Prostaglandin E₂ alleviates cyclosporin A-induced bone loss in the rat. *J. Bone and Min. Res.*, 7:1191-1200.
13. Ke, HZ, Li, M, Jee, WSS. 1992 Prostaglandin E₂ prevents ovariectomy-induced cancellous bone loss in rats. *Bone and Mineral* 19:45-62.

W.S.S. Jee, Ph.D.
NAG-24354

14. Ke, HZ, Jee, WSS, Mori, S, Li, XJ, Kimmel, DB. 1992 Effects of long-term daily administration of prostaglandin-E₂ on maintaining elevated proximal tibial metaphyseal cancellous bone mass in male rats. *Calcif. Tissue Int.* 50:245-252.
15. Ke, HZ, Jee, WSS. 1992 Effects of daily administration of prostaglandin E₂ and its withdrawal on the lumbar vertebral bodies in male rats. *Anat. Rec.* 234:172-182.
16. Mori, S, Jee, WSS, Li, XJ. 1992 Production of new trabecular bone in osteopenic ovariectomized rats by prostaglandin E₂. *Calcif. Tissue Int.* 50:80-87.
17. Tang, LY, Jee, WSS, Ke, HZ, Kimmel DB. 1992 Restoring and maintaining bone in osteopenic female rat skeleton: I. Changes in bone mass and structure. *J. Bone and Min. Res.* 7:1093-1104.
18. Ito, H., Ke, HZ, Jee, WSS, Sakou, T. 1993 Anabolic responses of an adult cancellous bone site to prostaglandin E₂ in the rat. *Bone and Mineral* 21:219-236.
19. Jee, WSS, Tang, L, Ke, HZ, Setterberg, RB, Kimmel, DB. 1993 Maintaining restored bone with bisphosphonate in the ovariectomized rat skeleton: Dynamic histomorphometry of changes in bone mass. *Bone* 14:493-498.
20. Ke, HZ, Jee, WSS, Zeng, QQ, Li, M, Lin BY. 1993 Prostaglandin E₂ increased rat cortical bone mass when administered immediately following ovariectomy. *Bone and Mineral* 21:189-201.
21. Ke, HZ, Jee, WSS, Ito, H, Setterberg, RB, Li, M, Lin, BY, Liang XG, Ma, YF. 1993 Greater bone formation induction occurred in aged than young cancellous bone sites. *Bone* 14:481-485.
22. Li, M, Jee, WSS, Ke, HZ, Liang, XG, Lin, BY, Ma, YF, Setterberg, RB. 1993 Prostaglandin E₂ restores cancellous bone to immobilized limb and adds bone to overloaded limb in right hindlimb immobilization rats. *Bone* 14:283-288.
23. Zeng, QQ, Jee, WSS, Ke, HZ, Wechter, WJ. 1993 S-Ketoprofen inhibits tenotomy-induced bone loss and dynamics in weanling rats. *Bone and Mineral* 21:203-218.
24. Frost, HM, Jee, WSS. 1994 Perspectives: A vital biomechanical model of the endochondral ossification mechanism. *Anat. Rec.* (In press).
25. Frost, HM, Jee, WSS. 1994 Perspectives: Applications of a biomechanical model of the endochondral ossification mechanism. *Anat. Rec.* (In press).

W.S.S. Jee, Ph.D.
NAG-2435

26. Jee, WSS, Ma, YF, Li, M. Liang, XG, Lin, BY, Li, XJ, Ke, HZ, Mori, S, Setterberg, RB, Kimmel, DB. 1994 Sex steroids and prostaglandins in bone metabolism. In: Sex Steroids and Bone. R. Ziegler, J. Pfeilschifter, M. Brautigam, eds. Springer-Verlag, Berlin. Pp. 119-150.
27. Lin, BY, Jee, WSS, Ma, YF, Ke, HZ, Kimmel, DB, Li, XJ. 1994 Effects of prostaglandin E₂ and risedronate administration on cancellous bone in older female rats. Bone 15:489-496.
28. Ma, YF, Ke, HZ, Jee, WSS. 1994 Prostaglandin E₂ adds bone to a cancellous bone site with a closed growth plate and low bone turnover in ovariectomized rats. Bone 15:137-146.
29. Ma, YF, Jee, WSS, Ke, HZ, Lin, BY, Liang, XG, Li, M, Yamamoto, N. 1994 Human parathyroid hormone (1-38) restores cancellous bone to the immobilized, osteopenic proximal tibial metaphysis in rats. (In press).29